arXiv:1207.7154v2 [math.DS] 3 Aug 2012

PATTERN GENERATION PROBLEMS ARISING IN MULTIPLICATIVE INTEGER SYSTEMS

JUNG-CHAO BAN*, WEN-GUEI HU**, AND SONG-SUN LIN†

Abstract. This study investigates a multiplicative integer system using a method that was developed for studying pattern generation problems. The entropy and the Minkowski dimensions of general multiplicative systems can thus be computed. A multi-dimensional decoupled system is investigated in three main steps. (I) Identify the admissible lattices of the system; (II) compute the density of copies of admissible lattices of the same length, and (III) compute the number of admissible patterns on the admissible lattices.

A coupled system can be decoupled by removing the multiplicative relation set and then performing procedures similar to those applied to a decoupled system . The admissible lattices are chosen to be the maximum graphs of different degrees which are mutually independent. The entropy can be obtained after the remaining error term is shown to approach zero as the degree of the admissible lattice tends to infinity.

1. Introduction

Multiplicative integer systems have been intensively studied in recent years; see [8, 9, 14, 17, 18] and the references therein. One of the main related issues is to compute Minkowski (box) dimension and Hausdorff dimension of such systems and to compare them. These two dimensions are equal in an additive shift. However, for most known examples of multiplicative integer system, they are different. Since the computations of these two dimensions are difficult, effective methods for computing these dimensions for general multiplicative systems must be develpoed.

This investigation is motivated directly by the work of Kenyon et al. [14], who utilized a variational method to obtain the results on

$$\mathbb{X}_2^0=\left\{(x_1,x_2,x_3,\cdots)\in\{0,1\}^{\mathbb{N}}\mid x_kx_{2k}=0 \text{ for all } k\geq 1\right\}, \tag{1.1}$$

and also pointed out that the method fails for the system

$$\mathbb{X}_{2,3}^0=\left\{(x_1,x_2,x_3,\cdots)\in\{0,1\}^{\mathbb{N}}\mid x_kx_{2k}x_{3k}=0 \text{ for all } k\geq 1\right\}. \tag{1.2}$$

This work provides an approach to general multiplicative systems, including (1.1) and (1.2).

---

*The author would like to thank the National Science Council, R.O.C. (Contract No. NSC 100-2115-M-259-009-MY2) and the National Center for Theoretical Sciences for partially supporting this research.

**The author would like to thank the National Science Council, R.O.C. and the ST Yau Center for partially supporting this research.

†The author would like to thank the National Science Council, R.O.C. (Contract No. NSC 98-2115-M-009) and the ST Yau Center for partially supporting this research.

This study emphasizes the computation of entropy $h(\mathbb{X})$ of the multiplicative system $\mathbb{X}$. The Minkowski dimension $\dim_M(\mathbb{X})$ is given by

$$\dim_M(\mathbb{X}) = \frac{1}{\log N} h(\mathbb{X}), \tag{1.3}$$

where N is the number of the symbols of system $\mathbb{X}$. The results of Hausdorff dimensions will appear elsewhere.

Previously, the authors studied pattern generation problems on multi-dimensional shifts (of finite type) and developed some efficient means of studying the generation of admissible patterns, and then computing the topological entropy (or spatial entropy); see [1, 2, 3, 4, 5, 6, 7, 10, 11, 12, 13, 15]. This study shows that these methods can be used to study the "multi-dimensional or coupled systems of multiplicative integers", including $\mathbb{X}^0_{2,3}$.

To illustrate our method, (1.1) is investigated first. The topological entropy $h(\mathbb{X}^0_2)$ has been shown to be

$$h(\mathbb{X}^0_2) = \sum_{k=1}^{\infty} \frac{1}{2^{k+1}} \log a_k, \tag{1.4}$$

where a_k is a Fibonacci number with $a_1 = 2$, $a_2 = 3$ and $a_{k+1} = a_k + a_{k-1}$ for all $k \geq 2$ [8]. The derivation of (1.4) is as follows. Denote by the (multiplicative) relation set $\mathbb{M}_2$ of the integers of 2-power, i.e.,

$$\mathbb{M}_2 = \{1, 2, 4, 8, 16, 32, \cdots, 2^n, \cdots\}. \tag{1.5}$$

Denote by the complementary index set $\mathcal{I}_2$ of $\mathbb{X}^0_2$ that contains all positive odd integers:

$$\mathcal{I}_2 = \{n \in \mathbb{N} \mid 2 \nmid n\} = \{2k+1\}_{k=0}^{\infty}. \tag{1.6}$$

The set of all natural number $\mathbb{N}$ can now be rearranged into

$$\mathbb{N} = \bigcup_{i \in \mathcal{I}_2} i\mathbb{M}_2, \tag{1.7}$$

where $i\mathbb{M}_2 = \left\{i, 2i, 2^2 i, \cdots, 2^n i, \cdots\right\}$. Clearly,

$$i\mathbb{M}_2 \cap j\mathbb{M}_2 = \emptyset \tag{1.8}$$

if $i, j \in \mathcal{I}_2$ and $i \neq j$. More precisely, the right-hand side of (1.7) can be expressed as an $\mathbb{N} \times \mathbb{N}$ table, Table 1.1.



Table 1.1.

For $k \geq 1$, let Z_k be the blank lattice of k cells in $\mathbb{Z}^1$, $k \geq 1$, as present in Fig. 1.1.



Z_k

Figure 1.1.

Let M_k and iM_k be the numbered lattices of the first k elements in $\mathbb{M}_2$ and in $i\mathbb{M}_2$ on the Z_k, respectively.



M_k iM_k

Figure 1.2.

Let

$$\mathcal{N}(m) = \{k \in \mathbb{N} \mid 1 \leq k \leq m\}, \tag{1.9}$$

be the set of natural numbers that are less than or equal to m. For each $n \geq 1$ and $1 \leq i \leq 2^n$, let

$$k_n(i) = \max\left\{k \mid i2^k \leq 2^n\right\} = \log_2 \left\lfloor \frac{2^n}{i} \right\rfloor, \tag{1.10}$$

where $\lfloor x \rfloor$ is the largest integer that is less than or equal to x.

Then, from Table 1.1, it is clear that

$$\mathcal{N}(2^n) = \bigcup_{i \in \mathcal{I}_2, 1 \leq i \leq 2^n} iM_{k_n(i)}. \tag{1.11}$$

For example, for $n = 4$, from Table 1.1,

$$\mathcal{N}(2^4) = M_5 \cup (3M_3) \cup (5M_2) \cup (7M_2) \cup (9M_1) \cup (11M_1) \cup (13M_1) \cup (15M_1). \tag{1.12}$$

In terms of blank lattices, the numbers in $\mathcal{N}(2^4)$ lie on

$$\text{one copy of } Z_5, \text{one copy of } Z_3, \text{two copies of } Z_2 \text{ and } 2^2 \text{ copies of } Z_1. \tag{1.13}$$

The result of general $\mathcal{N}(2^n)$ follows from the following proposition, which can be easily proven by mathematical induction.

Proposition 1.1. *For integers $Q \geq 2$ and $n \geq 1$,*

$$Q^n = (n+1) + n(Q-2) + (Q-1)^2 \sum_{k=1}^{n-1} k Q^{n-1-k}. \tag{1.14}$$

In particular,

$$2^n = (n+1) + \sum_{k=1}^{n-1} k 2^{n-1-k}. \tag{1.15}$$

Therefore, (1.15) states that the numbers in $\mathcal{N}(2^n)$ are spread out on blank lattices with one copy of Z_{n+1} and 2^{n-1-k} copies of Z_k, $1 \leq k \leq n-1$. In particular, setting $n = 4$ in (1.15) yields (1.13).

Now, consider again system $\mathbb{X}_2^0$ and the target formula (1.4). For any $m \geq 1$, let X_m be the set of all admissible m-sequences in $\mathbb{X}_2^0$:

$$X_m = \left\{ (x_1, x_2, \cdots, x_m) \in \{0,1\}^{\mathbb{Z}_m} \;\middle|\; x_k x_{2k} = 0 \text{ for all } k \geq 1 \text{ and } 2k \leq m \right\}. \tag{1.16}$$

Our purpose is to compute $|X_m|$, which is the number of elements in (1.16). The entropy $h(\mathbb{X}_2^0)$ follows from

$$h(\mathbb{X}_2^0) = \lim_{m \to \infty} \frac{1}{m} \log |X_m|. \tag{1.17}$$

The constraint

$$x_k x_{2k} = 0 \tag{1.18}$$

in (1.16) is the admissible condition of golden-mean shift on $\mathbb{Z}^1$, and it states that symbol 1 is not allowed immediately to follow symbol 1. Then, the forbidden set on Z_2 is $\{\boxed{1}\boxed{1}\}$. The transition matrix

$$G = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}. \tag{1.19}$$

Le Σ_k be the set of all admissible patterns on $\mathbb{Z}_k$ with respect to (1.19); then

$$|\Sigma_k| = a_k, \tag{1.20}$$

which is the k-th Fibonacci number. Since the constraint (1.18) applies to each $i\mathbb{M}_2$ independently for $i \in \mathcal{I}_2$,

$$|X_{2^n}| = |\Sigma_{n+1}| \prod_{k=1}^{n-1} |\Sigma_k|^{2^{n-1-k}}, \tag{1.21}$$

which implies

$$\frac{1}{2^n} \log |X_{2^n}| = \frac{1}{2^n} a_{n+1} + \sum_{k=1}^{n-1} \frac{1}{2^{k+1}} \log a_k. \tag{1.22}$$

Hence (1.4) follows easily from (1.22).

Clearly, (1.14) of Proposition 1.1 also recovers the following results [8].

Theorem 1.2. *For any $Q \geq 2$, denote the multiplicative integers system*

$$\mathbb{X}_Q^0 = \left\{ (x_1, x_2, \cdots) \in \{0,1\}^{\mathbb{N}} \mid x_k x_{Qk} = 0 \text{ for all } k \geq 1 \right\}, \tag{1.23}$$

then

$$h(\mathbb{X}_Q^0) = (Q-1)^2 \sum_{k=1}^{\infty} \frac{1}{Q^{k+1}} \log a_k. \tag{1.24}$$

Consideration of the above reveals the following three main parts of our study of $\mathbb{X}_2^0$.

(I) Identify the numbered lattice M_k and the "admissible" blank lattice Z_k from the given system; see Figs. 1.1 and 1.2.

(II) Compute the numbers of copies of independent admissible lattices of the same length; see formulae (1.14) and (1.15).

(III) Determine the set of all admissible patterns Σ_k, which can be generated on Z_k, and compute the number thereof $|\Sigma_k|$.

Notably, step (III) in the study of $\mathbb{X}_2^0$ is the classical one-dimensional pattern generation problem; see [16].

Based on the above observations, the rest of this paper will consider the following two classes of systems.

(i) Multi-dimensional systems like

$$\mathbb{X}_{2,3}^0 = \left\{ (x_1, x_2, x_3, \cdots) \in \{0,1\}^{\mathbb{N}} \mid x_k x_{2k} x_{3k} = 0 \text{ for all } k \geq 1 \right\}. \tag{1.25}$$

(ii) One-dimensional coupled systems like

$$\mathbb{X}_2^A = \{ (x_1, x_2, \cdots) \in \Sigma_A \mid x_k x_{2k} = 0 \text{ for all } k \geq 1 \}, \tag{1.26}$$

i.e.,

$$\mathbb{X}_2^A = \mathbb{X}_2^0 \cap \Sigma_A, \tag{1.27}$$

where Σ_A is a proper additive shift of finite type.

For the multi-dimensional coupled system like

$$\begin{aligned}\mathbb{X}^A_{2,3} = &\ \{(x_1,x_2,\cdots)\in\Sigma_A \mid x_k x_{2k} x_{3k}=0 \text{ for all } k\geq 1\} \\ = &\ \mathbb{X}^0_{2,3}\cap\Sigma_A,\end{aligned} \tag{1.28}$$

which is more delicate, will be studied in the consecutive paper.

After we completed our study of (i) and (ii), we became aware of the announcement of the work of Peres et al. [17] on (1.25). These authors obtained the same results as ours for multi-dimensional system (i). Our methods for studying (i) differ from theirs. Moreover, our methods can be applied to (ii). To illustrate these methods and also for completeness, our approach to (i) is presented as involving the three specified steps (I), (II) and (III). Later, a modification of these procedures enables us to study the coupled system.

The rest of this paper is arranged as follows. Section 2 studies multi-dimensional systems. Section 3 studies coupled systems.

2. Multi-dimensional systems

This section concerns multi-dimensional systems. For simplicity, $\mathbb{X}^0_{2,3}$ is considered first. Recall that

$$\mathbb{X}^0_{2,3}=\left\{(x_1,x_2,x_3,\cdots)\in\{0,1\}^{\mathbb{N}} \mid x_k x_{2k} x_{3k}=0 \text{ for all } k\geq 1\right\}. \tag{2.1}$$

Firstly, the admissible numbered and blank lattices, determined by the constraint

$$x_k x_{2k} x_{3k}=0, \tag{2.2}$$

must be identified in $\mathbb{X}^0_{2,3}$.

Let $\mathbb{M}_{2,3}$ be the set of the numbers that are multiples of 2-power and 3-power:

$$\mathbb{M}_{2,3}=\left\{2^k 3^l \ \mid\ k,l\geq 0\right\}. \tag{2.3}$$

$\mathbb{M}_{2,3}$ can be expressed in the first quadrant of the plane, as follows.

243	486	972	1944	3888	7776
81	162	324	648	1296	2592
27	54	108	216	432	864
9	18	36	72	144	288
3	6	12	24	48	96
1	2	4	8	16	32

q_{27}	q_{33}	q_{40}	q_{47}	q_{55}	q_{64}
q_{19}	q_{24}	q_{30}	q_{36}	q_{43}	q_{51}
q_{12}	q_{16}	q_{21}	q_{26}	q_{32}	q_{39}
q_7	q_{10}	q_{14}	q_{18}	q_{23}	q_{29}
q_3	q_5	q_8	q_{11}	q_{15}	q_{20}
q_1	q_2	q_4	q_6	q_9	q_{13}

$\mathbb{M}_{2,3}$

Table 2.1.

Since $\mathbb{M}_{2,3}$ inherits the natural ordering of natural numbers, the k-th number in $\mathbb{M}_{2,3}$ can be denoted by q_k, $k\in\mathbb{N}$. It seems that no known formula relates $2^m 3^n$ to q_i explicitly [19].

From Table 2.1, for any $k\geq 1$, the L-shaped k-cell numbered lattice M_k that contains $\{q_1,q_2,\cdots,q_k\}$ can be defined, and k-cell blank lattice L_k can also be defined by deleting the numbers of M_k. Table 2.2 presents the first 15 numbered lattices M_k. Notably, the number q_k in the bold square is the largest number in M_k.



Table 2.2.

In contrast to $\mathbb{X}_2^0$, all lattices M_k and L_k are now two-dimensional. Therefore, the system $\mathbb{X}_{2,3}^0$ can be regarded as a two-dimensional system; see [3, 5, 6, 7, 15].

Now, the complementary index set $\mathcal{I}_{2,3}$ of $\mathbb{X}_{2,3}^0$ is the set of all natural numbers that cannot be divided by two and three:

$$\begin{aligned}\mathcal{I}_{2,3} = \ & \{n \in \mathbb{N} \ | \ 2 \nmid n \text{ and } 3 \nmid n\} \\ = \ & \{6k+1, 6k+5\}_{k=0}^{\infty} \\ = \ & \{1, 5, 7, 11, 13, 17, 19, 23, 25, 29, \cdots\}.\end{aligned} \tag{2.4}$$

The set $\mathbb{N}$ of natural numbers can be rearranged into the first octant of three-dimensional space as

$$\mathbb{N} = \bigcup_{i \in \mathcal{I}_{2,3}} i\mathbb{M}_{2,3}, \tag{2.5}$$

and for $i, j \in \mathcal{I}_{2,3}$ with $i \neq j$,

$$i\mathbb{M}_{2,3} \cap j\mathbb{M}_{2,3} = \emptyset; \tag{2.6}$$

see Fig. 2.1.



Figure 2.1.

Therefore, for any $n \in \mathbb{N}$, there exists a unique $i = i(n)$ such that $n \in i\mathbb{M}_{2,3}$.

Next, proceed to step (II): compute the numbers of copies of M_k for a given $\mathcal{N}(m)$, defined in (1.9).

Let

$$q_K = 2^m 3^n \in \mathbb{M}_{2,3} \tag{2.7}$$

for some $K \geq 1$. Denote by

$$J_K = \{ i \in \mathcal{I}_{2,3} \mid 1 \leq i \leq q_K \} . \tag{2.8}$$

Define

$$q_K(i) = \max \{ q \in \mathbb{M}_{2,3} \mid iq \leq q_K \} , \tag{2.9}$$

then $q_K(i) \geq 1$ for any $i \in J_K$.

Denote by the set

$$I_K(k) = \{ i \in J_K \mid q_K(i) = k \} \tag{2.10}$$

for $1 \leq k \leq K$. Clearly, if $q_K(i) = k$, then

$$iq_k \leq q_K < iq_{k+1}.$$

Therefore, a parallel result of (1.11) and (1.14) is as follows.

$$\mathcal{N}(q_K) = \bigcup_{i \in J_K} iM_{q_K(i)} = \bigcup_{i \in I_K(k)} iM_k \tag{2.11}$$

with

$$I_K(k) = \left(\frac{q_K}{q_{k+1}}, \frac{q_K}{q_k} \right] \cap \mathcal{I}_{2,3}. \tag{2.12}$$

The following example illustrates (2.11) and (2.12).

Example 2.1. *For* $q_{14} = 36$, *it is easy to verify*

$$\begin{cases} I_{14}(1) = \{19, 23, 25, 29, 31, 35\} \\ I_{14}(2) = \{13, 17\} \\ I_{14}(3) = \{11\}, \ \ I_{14}(4) = \{7\}, \ \ I_{14}(5) = \{5\}, \ \ I_{14}(14) = \{1\}. \end{cases}$$

The others are empty. Therefore,

$$\mathcal{N}(36) = M_{14} \cup 5M_5 \cup 7M_4 \cup 11M_3 \cup 13M_2 \cup 17M_2 \bigcup_{i\in\{19,23,25,29,31,35\}} iM_1.$$

For $1 \le k \le K$, denote by

$$\alpha_K(k) = |I_K(k)|, \tag{2.13}$$

the number of copies of M_k in $\mathcal{N}(q_K)$.

Formula (2.12) gives the density of copies of M_k in $[1, q_K]$, which is crucial in computing the entropy. Indeed, following proposition is offered.

Proposition 2.2. *On* $\mathbb{X}^0_{2,3}$, *for an* $k \ge 1$,

$$\lim_{K\to\infty} \frac{\alpha_K(k)}{q_K} = \beta_{2,3}\left(\frac{1}{q_k} - \frac{1}{q_{k+1}}\right), \tag{2.14}$$

where

$$\beta_{2,3} = \frac{\sharp\left\{\mathcal{I}_{2,3} \cap [1, [2,3]]\right\}}{[2,3]} = \frac{1}{3}, \tag{2.15}$$

and $[2, 3] = 6$ *is the least common multiple of* 2 *and* 3*.*

Proof. For any fixed $k \ge 1$, from (2.12) and (2.13), (2.14) follows. Clearly, (2.15) follows from (2.4). □

It remains to show that limit (2.14) exists not only for q_K but for all n tends to infinity. More precisely, for any $1 < l \in \mathcal{I}_{2,3}$, denote by

$$\begin{cases} \mathcal{N}_{K,l} = \left\{n \in \mathbb{N} \ \mid \ 1 \le n \le lq_K\right\}, \\ J_{K,l} = \left\{i \in \mathcal{I}_{2,3} \ \mid \ i \le lq_K\right\}, \\ q_{K,l}(i) = \max\left\{q \in \mathbb{M}_{2,3} \ \mid \ iq \le lq_K\right\}, \\ I_{K,l}(k) = \left\{i \in J_{K,l} \ \mid \ q_{K,l}(i) = k\right\} \text{ and} \\ \alpha_{K,l}(k) = |I_{K,l}(k)|. \end{cases} \tag{2.16}$$

The following proposition can be easily verified. The detailed proof is omitted.

Proposition 2.3. *For* $k \ge 1$,

$$\lim_{K\to\infty} \frac{\alpha_{K,l}(k)}{lq_K} = \beta_{2,3}\left(\frac{1}{q_k} - \frac{1}{q_{k+1}}\right), \tag{2.17}$$

where $\beta_{2,3}$ *is given by (2.15).*

After the density of M_k (2.14) is obtained, step (II) is completed. Now, the final step (III) is to compute the admissible patterns on L_k for all $k \geq 1$.

Previously, two-dimensional pattern generation problems on L-shaped lattices has been studied by Lin and Yang in [15]. The constraint (2.2) implies that the forbidden local patterns on L_3 are

$$\mathcal{F}_{2,3} = \left\{ \begin{array}{cc} \boxed{1} & \\ \boxed{1} & \boxed{1} \end{array} \right\}. \tag{2.18}$$

Therefore, the basic set of admissible patterns is

$$\Sigma_3 = \mathcal{B}_{2,3} = \left\{ \begin{array}{cc} \boxed{0} & \\ \boxed{0} & \boxed{0} \end{array}, \begin{array}{cc} \boxed{1} & \\ \boxed{0} & \boxed{0} \end{array}, \begin{array}{cc} \boxed{0} & \\ \boxed{0} & \boxed{1} \end{array}, \begin{array}{cc} \boxed{1} & \\ \boxed{0} & \boxed{1} \end{array}, \begin{array}{cc} \boxed{0} & \\ \boxed{1} & \boxed{0} \end{array}, \begin{array}{cc} \boxed{1} & \\ \boxed{1} & \boxed{0} \end{array}, \begin{array}{cc} \boxed{0} & \\ \boxed{1} & \boxed{1} \end{array} \right\}; \tag{2.19}$$

see [5].

Clearly, $\Sigma_1 = \left\{ \boxed{0}, \ \boxed{1} \right\}$ and $\Sigma_2 = \left\{ \boxed{0\,|\,0}, \ \boxed{0\,|\,1}, \ \boxed{1\,|\,0}, \ \boxed{1\,|\,1} \right\}$, and no constraint applies on L_1 and L_2. For $k \geq 4$, L_k can be regarded as the union of L_2 and L_3, and the imposed constraint only applies on L_3 parts. Hence, for $k \geq 4$, the admissible patterns on L_k are completely determined by $\mathcal{B}_{2,3}$.

After the above procedures have been completed, the following result concerning the entropy $h(\mathbb{X}^0_{2,3})$ is obtained.

Theorem 2.4. *The entropy of* $\mathbb{X}^0_{2,3}$ *is given by*

$$h(\mathbb{X}^0_{2,3}) = \sum_{k=1}^{\infty} \beta_{2,3} \left(\frac{1}{q_k} - \frac{1}{q_{k+1}} \right) \log |\Sigma_k|, \tag{2.20}$$

where $|\Sigma_k|$ *is the number of all admissible patterns determined by* $\mathcal{B}_{2,3}$.

Proof. The proof follows from Propositions 2.2 and 2.3. □

Denote by

$$\Sigma_k = \Sigma_k(\mathcal{B}_{2,3}) \tag{2.21}$$

the set of all admissible patterns that are determined by $\mathcal{B}_{2,3}$. Denote by

$$|\Sigma_k| = b_k \tag{2.22}$$

the number of patterns in Σ_k. Since no exact formula relates $2^m 3^n$ to q_i for $\mathbb{M}_{2,3}$ in Table 2.1, unlike for a Fibonacci number, no "recursive" formula exists for b_k. This fact creates the most difficulty in computing entropy for a multi-dimensional system; see[3, 5, 6, 7, 15].

However, for relatively small k, b_k can be computed. Table 2.3 presents cases for $k \in [1, 14]$ and k with $q_{k+1} = 6^l$, $1 \leq l \leq 8$. Clearly, $q_5 = 6$, $q_{14} = 6^2$, $q_{26} = 6^3$, $q_{43} = 6^4$, $q_{64} = 6^5$, $q_{89} = 6^6$, $q_{119} = 6^7$, $q_{153} = 6^8$. Remark 2.5 concerns a further property of b_k.

k	1	2	3	4	5	6	7	8	9	10	11	12	13	14
b_k	2	4	7	14	25	50	90	160	320	584	1039	1861	3722	6774

$b_{25} = 5,434,757$	$b_{42} = 172,749,984,030$
$b_{63} \approx 5.291646495998910 \times 10^{16}$	$b_{88} \approx 2.006283543836154 \times 10^{23}$
$b_{118} \approx 1.439075072036499 \times 10^{31}$	$b_{152} \approx 1.022234008604671 \times 10^{40}$

Table 2.3.

In Table 2.3, b_k has been computed explicitly up to $k = 152$ by using the transition matrices developed in [5, 6].

Remark 2.5. *Define the ratio of* $|\Sigma_k|$ *by*

$$r_k = |\Sigma_k|/|\Sigma_{k-1}| = b_k/b_{k-1} \tag{2.23}$$

for $k \geq 2$. *From Tables 2.1, 2.2 and 2.3, it is easy to verify that*

$$r_k = 2 \quad \text{if} \quad q_k = 2^n, \tag{2.24}$$

for some $n \geq 1$.

On the other hand, it can be shown that there exists $C \leq \frac{31}{16}$ *such that*

$$r_k \leq C \quad \text{for} \quad q_k \neq 2^n \text{ for all } n \geq 2. \tag{2.25}$$

Therefore, $\{r_k\}$ *cannot have a limit as* k *tends to* ∞, *unlike the Fibonacci sequence, which has the limit* $\frac{1+\sqrt{5}}{2}$. *A detailed study of* $\{r_k\}$ *and* b_k *will appear elsewhere.*

In the following, an approximation of (2.20) is given. For $n \geq 1$, let

$$h^{(n)}(\mathbb{X}^0_{2,3}) = \sum_{k=1}^{n} \beta_{2,3} \left(\frac{1}{q_k} - \frac{1}{q_{k+1}} \right) \log |\Sigma_k|. \tag{2.26}$$

Clearly, from Theorem 2.4, $h^{(n)}(\mathbb{X}^0_{2,3})$ is a lower bound of $h(\mathbb{X}^0_{2,3})$, and $h^{(n)}(\mathbb{X}^0_{2,3})$ increasingly approaches $h(\mathbb{X}^0_{2,3})$ as n tends to infinity. Furthermore, let

$$\begin{aligned} E^{(n)}(\mathbb{X}^0_{2,3}) \equiv & \sum_{k=n+1}^{\infty} \beta_{2,3} \left(\tfrac{1}{q_k} - \tfrac{1}{q_{k+1}} \right) \log(2^k) \\ = & \ \beta_{2,3} \log 2 \left(\tfrac{n+1}{q_{n+1}} + 3 - \sum_{k=1}^{n+1} \tfrac{1}{q_k} \right), \end{aligned} \tag{2.27}$$

where $\sum\limits_{k=1}^{\infty} \frac{1}{q_k} = 3$. Hence,

$$h(\mathbb{X}^0_{2,3}) - h^{(n)}(\mathbb{X}^0_{2,3}) \leq E^{(n)}(\mathbb{X}^0_{2,3}). \tag{2.28}$$

Table 2.4 presents cases for n with $q_{n+1} = 6^l$ and $1 \leq l \leq 8$.

n	4	13	25	42	63	88	118	152
$h^{(n)}(\mathbb{X}^0_{2,3})$	0.319901	0.537229	0.620707	0.645733	0.652284	0.653865	0.654224	0.654303

Table 2.4.

Moreover, $E^{(152)}(\mathbb{X}^0_{2,3}) \approx 0.0000238741$.

The theorem just established can be easily extended to general multiplicative systems. For completeness, the result is simply stated and necessary modifications to its proof are only sketched.

Consider any two natural numbers $\gamma_1 < \gamma_2$ such that γ_2 is not a pure power of γ_1:

$$\gamma_2 \neq \gamma_1^m \tag{2.29}$$

for all $m \geq 1$. Let

$$\mathbb{X}^0_{\gamma_1,\gamma_2} = \left\{(x_1, x_2, x_3, \cdots) \in \{0,1\}^{\mathbb{N}} \mid x_k x_{\gamma_1 k} x_{\gamma_2 k} = 0 \text{ for all } k \geq 1\right\}. \tag{2.30}$$

Denote by

$$\begin{aligned} \mathbb{M}_{\gamma_1,\gamma_2} = & \quad \{\gamma_1^m \gamma_2^n \mid m, n \geq 0\} \\ = & \quad \{q_1, q_2, q_3, \cdots\}, \end{aligned} \tag{2.31}$$

on the first quadrant on the plane as Tables 1.1 and 2.1, where

$$q_k = q_k(\gamma_1, \gamma_2). \tag{2.32}$$

Then, the numbered lattice $M_k = M_k(\gamma_1, \gamma_2)$ and the blank lattice $L_k = L_k(\gamma_1, \gamma_2)$ can be obtained from (2.31) for $k \geq 1$.

The complementary index set

$$\mathcal{I}_{\gamma_1,\gamma_2} = \{n \in \mathbb{N} \mid \gamma_1 \nmid n \text{ and } \gamma_2 \nmid n\}. \tag{2.33}$$

Then

$$\mathbb{N} = \bigcup_{i \in \mathcal{I}_{\gamma_1,\gamma_2}} i\mathbb{M}_{\gamma_1,\gamma_2}. \tag{2.34}$$

Let $\Sigma_k(\gamma_1, \gamma_2)$ be the set of admissible patterns on L_k. Then, the entropy of $\mathbb{X}^0_{\gamma_1,\gamma_2}$ is given by the following theorem.

Theorem 2.6. *For any two integer numbers $\gamma_2 > \gamma_1 > 1$ with $\gamma_2 \neq \gamma_1^m$ for all $m > 1$. Then,*

$$h(\mathbb{X}^0_{\gamma_1,\gamma_2}) = \sum_{k=1}^{\infty} \beta_{\gamma_1,\gamma_2} \left(\frac{1}{q_k} - \frac{1}{q_{k+1}}\right) \log |\Sigma_k(\gamma_1, \gamma_2)|, \tag{2.35}$$

where

$$\beta_{\gamma_1,\gamma_2} = \frac{\sharp\{\mathcal{I}_{\gamma_1,\gamma_2} \cap [1, [\gamma_1,\gamma_2]]\}}{[\gamma_1,\gamma_2]} \tag{2.36}$$

where $[\gamma_1,\gamma_2]$ *is the least common multiple of* γ_1 *and* γ_2.

The following example illustrates Theorem 2.8.

Example 2.7. *Consider* $\mathbb{X}^0_{2,5}$. *Then*

625	1250	2500	5000	10000	20000
125	250	500	1000	2000	4000
25	50	100	200	400	800
5	10	20	40	80	160
1	2	4	8	16	32

q_{26}	q_{31}	q_{36}	q_{42}	q_{48}	q_{55}
q_{16}	q_{20}	q_{24}	q_{29}	q_{34}	q_{40}
q_9	q_{12}	q_{15}	q_{19}	q_{23}	q_{28}
q_4	q_6	q_8	q_{11}	q_{14}	q_{18}
q_1	q_2	q_3	q_5	q_7	q_{10}

$\mathbb{M}_{2,5}$

Table 2.5.

Notably, starting from $k \geq 5$, *all* L_k *in* $\mathbb{M}_{2,5}$ *differ from those in* $\mathbb{M}_{2,3}$, *so* $\mathbb{X}^0_{2,3}$ *and* $\mathbb{X}^0_{2,5}$ *have different sequences of admissible blank lattices, affecting* $\Sigma_k(\gamma_1,\gamma_2)$. *For example,*



Table 2.6.

Table 2.7 presents cases for n *with* $q_{n+1} = 10^l$ *and* $1 \leq l \leq 7$. *Clearly,* $q_6 = 10$, $q_{15} = 10^2$, $q_{29} = 10^3$, $q_{48} = 10^4$, $q_{72} = 10^5$, $q_{100} = 10^6$, $q_{134} = 10^7$.

n	$\lvert\Sigma_n(2,5)\rvert$	$h^{(n)}(\mathbb{X}^0_{2,5})$
5	*28*	*0.431341*
14	*7,402*	*0.616889*
28	*38,739,020*	*0.660010*
47	*4,446,117,963,472*	*0.667813*
71	$\approx 8.805722546306860 \times 10^{18}$	*0.669042*
99	$\approx 1.961346392265600 \times 10^{26}$	*0.669220*
133	$\approx 1.685495616218491 \times 10^{35}$	*0.669244*

Table 2.7.

Moreover, $E^{(133)}(\mathbb{X}^0_{2,5}) \approx 0.0000259978$.

On the other hand, if

$$\gamma_1 = Q \quad \text{and} \quad \gamma_2 = Q^m, \tag{2.37}$$

where $Q \geq 2$ and $m \geq 2$, then $\mathbb{X}^0_{Q,Q^m}$ is reduced to a one-dimensional problem and the following result is obtained.

Theorem 2.8. *For $Q, m \geq 2$, if $\gamma_1 = Q$ and $\gamma_2 = Q^m$, then*

$$h(\mathbb{X}^0_{Q,Q^m}) = (Q-1)^2 \sum_{k=1}^{\infty} \frac{1}{Q^{k+1}} \log |a_k(Q,Q^m)| \,, \tag{2.38}$$

where $a_k = |A(Q,Q^m)|$ for $k \geq m$, $a_j = Q^j$, $1 \leq j \leq m$, where $A(Q,Q^m)$ is the associated transition matrix of $\mathcal{B}(Q,Q^m)$.

The following two examples illustrate Theorem 2.8.

Example 2.9. *First, consider $\mathbb{X}^0_{2,4}$. Steps (I) and (II) are the same as for $\mathbb{X}^0_2$. In step (III), the forbidden set of $\Sigma_k(2,4)$ is $\{\boxed{\begin{array}{c|c|c}1&1&1\end{array}}\}$. The associated transition matrix*

$$A(2,4) = \begin{array}{c} \\ \boxed{\begin{array}{c|c}0&0\end{array}} \\ \boxed{\begin{array}{c|c}0&1\end{array}} \\ \boxed{\begin{array}{c|c}1&0\end{array}} \\ \boxed{\begin{array}{c|c}1&1\end{array}} \end{array} \begin{array}{c} \begin{array}{cccc} \boxed{\begin{array}{c|c}0&0\end{array}} & \boxed{\begin{array}{c|c}0&1\end{array}} & \boxed{\begin{array}{c|c}1&0\end{array}} & \boxed{\begin{array}{c|c}1&1\end{array}} \end{array} \\ \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \end{array} . \tag{2.39}$$

Therefore, as in (1.4),

$$h(\mathbb{X}^0_{2,4}) = \sum_{k=1}^{\infty} \frac{1}{2^{k+1}} \log a_k(2,4), \tag{2.40}$$

where $a_1(2,4) = 2$, $a_2(2,4) = 4$ and $a_k(2,4) = \left|A(2,4)^{k-2}\right|$ for all $k \geq 3$, where $|A|$ is the sum of all entries in A.

Next, consider $\mathbb{X}^0_{2,8}$. Now, the forbidden set of $\Sigma_k(2,8)$ is $\left\{\boxed{\begin{array}{c|c|c|c}1&1&0&1\end{array}}, \boxed{\begin{array}{c|c|c|c}1&1&1&1\end{array}}\right\}$. Hence, the associated transition matrix

$$A(2,8) = \begin{array}{c} \\ \boxed{\begin{array}{c|c|c}0&0&0\end{array}} \\ \boxed{\begin{array}{c|c|c}0&0&1\end{array}} \\ \boxed{\begin{array}{c|c|c}0&1&0\end{array}} \\ \boxed{\begin{array}{c|c|c}0&1&1\end{array}} \\ \boxed{\begin{array}{c|c|c}1&0&0\end{array}} \\ \boxed{\begin{array}{c|c|c}1&0&1\end{array}} \\ \boxed{\begin{array}{c|c|c}1&1&0\end{array}} \\ \boxed{\begin{array}{c|c|c}1&1&1\end{array}} \end{array} \begin{array}{c} \begin{array}{cccccccc} \boxed{\begin{array}{c|c|c}0&0&0\end{array}} & \boxed{\begin{array}{c|c|c}0&0&1\end{array}} & \boxed{\begin{array}{c|c|c}0&1&0\end{array}} & \boxed{\begin{array}{c|c|c}0&1&1\end{array}} & \boxed{\begin{array}{c|c|c}1&0&0\end{array}} & \boxed{\begin{array}{c|c|c}1&0&1\end{array}} & \boxed{\begin{array}{c|c|c}1&1&0\end{array}} & \boxed{\begin{array}{c|c|c}1&1&1\end{array}} \end{array} \\ \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix} \end{array} .$$

Therefore,

$$h(\mathbb{X}^0_{2,8}) = \sum_{k=1}^{\infty} \frac{1}{2^{k+1}} \log a_k(2,8),$$

where $a_1(2,8) = 2$, $a_2(2,8) = 4$, $a_3(2,8) = 8$ and $a_k(2,8) = \left|A(2,8)^{k-3}\right|$ for all $k \geq 4$.

Now, the study of higher-dimensional systems is briefly described. The methods employed and results obtained for two-dimensional cases can easily be extended to higher-dimensional cases. Only the necessary modifications are sketched.

Let

$$1 < \gamma_1 < \gamma_2 < \cdots < \gamma_d \tag{2.41}$$

be natural numbers, $d \geq 3$, such that γ_j is not pure power of γ_i for all $i < j$, i.e.,

$$\gamma_j \neq \gamma_i^m, \tag{2.42}$$

for all $m \geq 2$ and $1 \leq i < j \leq d$.

Denote by

$$\Gamma \equiv \Gamma_d = (\gamma_1, \gamma_2, \cdots, \gamma_d) \tag{2.43}$$

and

$$\begin{aligned} \mathbb{X}_\Gamma^0 \equiv \; & \mathbb{X}^0_{\gamma_1, \gamma_2, \cdots, \gamma_d} \\ = \; & \left\{ (x_1, x_2, x_3, \cdots) \in \{0,1\}^{\mathbb{N}} \mid x_k x_{\gamma_1 k} x_{\gamma_2 k} \cdots x_{\gamma_d k} = 0 \text{ for all } k \geq 1 \right\} \end{aligned} \tag{2.44}$$

Denote by

$$\begin{aligned} \mathbb{M}_\Gamma \equiv \; & \{\gamma_1^{m_1} \gamma_2^{m_2} \cdots \gamma_d^{m_d} \;\mid\; m_j \geq 0\} \\ \equiv \; & \{q_k\}_{k=1}^{\infty} \end{aligned} \tag{2.45}$$

with $q_k < q_j$ if $k < j$.

Then, (2.45) defines a sequence of d-dimensional numbered lattices M_k of k cells. The blank lattices L_k are defined analogously. The complementary index set $\mathcal{I}_\Gamma$ of $\mathbb{M}_\Gamma$ is defined by

$$\mathcal{I}_\Gamma = \{ n \in \mathbb{N} \;\mid\; \gamma_j \nmid n, 1 \leq j \leq d \}. \tag{2.46}$$

Hence,

$$\mathbb{N} = \bigcup_{i \in \mathcal{I}_\Gamma} i \mathbb{M}_\Gamma. \tag{2.47}$$

The constraint

$$x_k x_{\gamma_1 k} x_{\gamma_2 k} \cdots x_{\gamma_d k} = 0 \tag{2.48}$$

determines the set $\mathcal{B}_\Gamma$ of basic admissible patterns which induces a d-dimensional shift of finite type $\Sigma(\mathcal{B}_\Gamma)$ on the d-dimensional lattice space $\mathbb{Z}^d$. Therefore, the admissible set $\Sigma_k(\mathcal{B}_\Gamma)$ is completely determined by $\mathcal{B}_\Gamma$ on L_k.

Now, by a similar argument, the entropy of $\mathbb{X}_\Gamma^0$ is easily obtained as follows. The detailed proof is omitted.

Theorem 2.10. *Let $\Gamma = \{\gamma_1, \gamma_2, \cdots, \gamma_d\}$ satisfy (2.41) and (2.42). Then the entropy of $\mathbb{X}_\Gamma^0$ is given by*

$$h(\mathbb{X}_\Gamma^0) = \sum_{k=1}^{\infty} \beta_\Gamma \left(\frac{1}{q_k} - \frac{1}{q_{k+1}} \right) \log |\Sigma_k|, \tag{2.49}$$

where

$$\beta_\Gamma = \frac{\mathcal{I}_\Gamma \cap [1,[\gamma_1,\gamma_2,\cdots,\gamma_d]]}{[\gamma_1,\gamma_2,\cdots,\gamma_d]} \tag{2.50}$$

and $[\gamma_1,\gamma_2,\cdots,\gamma_d]$ *is the least common multiple of* $\gamma_1,\gamma_2,\cdots,\gamma_d$.

The following three-dimensional system illustrates the methods and results.

Example 2.11. *For* $d=3$, *consider*

$$\mathbb{X}^0_{2,3,5} = \left\{(x_1,x_2,x_3,\cdots)\in\{0,1\}^{\mathbb{N}} \mid x_k x_{2k} x_{3k} x_{5k} = 0 \text{ for all } k\geq 1\right\}.$$

Then,



$\mathbb{M}_{2,3,5}$

Table 2.8

The first six numbered lattices are listed as follows.



Table 2.9

Clearly,

$$\mathcal{I}_{2,3,5} = \{30k+j \mid j\in\{1,7,11,13,17,19,23,29\} \text{ and } k\geq 0\}.$$

Therefore, it can be verified that

$$h(\mathbb{X}^0_{2,3,5}) = \sum_{k=1}^{\infty}\beta_{2,3,5}\left(\frac{1}{q_k}-\frac{1}{q_{k+1}}\right)\log|\Sigma_k(2,3,5)|, \tag{2.51}$$

where $\beta_{2,3,5}=\frac{4}{15}$ *and the forbidden set of* $\Sigma_k(2,3,5)$ *is* $\Bigg\{$  $\Bigg\}$.

The n-th order approximation of (2.51) is as follows. For $n \geq 1$, let

$$h^{(n)}(\mathbb{X}^0_{2,3,5}) = \sum_{k=1}^{n} \beta_{2,3,5} \left(\frac{1}{q_k} - \frac{1}{q_{k+1}} \right) \log |\Sigma_k(2,3,5)|.$$

In Table 2.10, $|\Sigma_n(2,3,5)|$, $1 \leq n \leq 25$, is listed.

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
$\lvert\Sigma_n(2,3,5)\rvert$	2	4	8	16	30	60	120	240	452	904	1708	3416	6832	12860	25720	48720	97440	184704	369408	738816	1390768	2625128	5250256	9979776	19959552

Table 2.10.

Moreover, $h^{(25)}(\mathbb{X}^0_{2,3,5}) \approx 0.548837$.

The constraint (2.48) is not essential and can be generalized. Therefore, the study of system $\mathbb{X}^0_\Gamma$ can also be generalized. In the following, the procedures are sketched below.

For any $N \geq 2$, consider a multiplicative system is of N-symbols, $\{0, 1, 2, \cdots, N-1\}$. For any $d \geq 1$, let the constraint set $\mathcal{C}$ be a subset of $\left\{0, 1, \cdots, (N-1)^d\right\}$. Denote by $\mathbb{X}_\Gamma(N, \mathcal{C})$ the multiplicative integer system with constraint set $\mathcal{C}$:

(2.52)
$$\mathbb{X}_\Gamma(N, \mathcal{C}) = \left\{ (x_1, x_2, x_3, \cdots) \in \{0, 1, \cdots, N-1\}^{\mathbb{N}} \mid x_k x_{\gamma_1 k} x_{\gamma_2 k} \cdots x_{\gamma_d k} \in \mathcal{C} \text{ for all } k \geq 1 \right\},$$

which consists of N-symbols, $\{0, 1, 2, \cdots, N-1\}$.

The following theorem can be easily obtained.

Theorem 2.12. *Let $\Gamma = \{\gamma_1, \gamma_2, \cdots, \gamma_d\}$ satisfy (2.41)∼(2.43) and $\mathcal{C} \subseteq \left\{0, 1, \cdots, (N-1)^d\right\}$. The entropy of $\mathbb{X}_\Gamma(N, \mathcal{C})$ is given by*

$$h(\mathbb{X}_\Gamma(N, \mathcal{C})) = \sum_{k=1}^{\infty} \beta_\Gamma \left(\frac{1}{q_k} - \frac{1}{q_{k+1}} \right) \log |\Sigma_k(\mathcal{B}_\Gamma(N, \mathcal{C}))|, \tag{2.53}$$

where $\Sigma_k(\mathcal{B}_\Gamma(N, \mathcal{C}))$ is the set of $d-$dimensional admissible local patterns that can be generated by $\mathcal{B}_\Gamma(N, \mathcal{C})$ on L_k.

The following example illustrates Theorem 2.12.

Example 2.13. *Let $N = 3$ and $\mathcal{C} = \{0, 2\}$. Then*

$$\mathbb{X}_2(3, \mathcal{C}) = \left\{ (x_1, x_2, x_3, \cdots) \in \{0, 1, 2\}^{\mathbb{N}} \mid x_k x_{2k} \in \{0, 2\} \textit{ for all } k \geq 1 \right\}.$$

The basic set of admissible local patterns is now given by

$$\mathcal{B}_2(3, \mathcal{C}) = \left\{ \ \boxed{0}\boxed{0}, \quad \boxed{0}\boxed{1}, \quad \boxed{0}\boxed{2}, \quad \boxed{1}\boxed{0}, \quad \boxed{1}\boxed{2}, \quad \boxed{2}\boxed{0}, \quad \boxed{2}\boxed{1} \ \right\}.$$

The associated transition matrix is

$$A(2;3,\mathcal{C})= \begin{array}{c} \\ \boxed{0} \\ \boxed{1} \\ \boxed{2} \end{array} \begin{array}{c} \begin{array}{ccc} \boxed{0} & \boxed{1} & \boxed{2} \end{array} \\ \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{array}\right] \end{array} .$$

Therefore, as in Theorem 1.2,

$$h(\mathbb{X}_2(3,\mathcal{C})) = 4 \sum_{k=1}^{\infty} \frac{1}{3^{k+1}} \log a_k(2;3,\mathcal{C}),$$

where $a_1(2;3,\mathcal{C}) = 3$*,* $a_k(2;3,\mathcal{C}) = \left|A(2;3,\mathcal{C})^{k-1}\right|$ *for all* $k \geq 2$*.*

The system (2.52) with $\gamma_j = \gamma_i^m$, $i < j$, for some $m \geq 2$ can also be studied. Detailed results are omitted.

3. Coupled systems

This section considers the coupled system, which is an intersection of the multiplicative integer systems $\mathbb{X}_Q^0$ with an additive proper subshift of finite type Σ_A,

$$\mathbb{X}_Q^A \equiv \mathbb{X}_Q^0 \cap \Sigma_A. \tag{3.1}$$

A simple system is considered first; the findings are then extended to general systems. Consider

$$\begin{aligned} \mathbb{X}_2^A \equiv\ & \mathbb{X}_2^0 \cap \Sigma_A \\ =\ & \left\{(x_1,x_2,x_3,\cdots) \in \{0,1\}^{\mathbb{N}} \mid x_k x_{2k} = 0 \text{ for all } k \geq 1 \text{ and } (x_1,x_2,x_3,\cdots) \in \Sigma_A\right\}, \end{aligned} \tag{3.2}$$

where

$$A = G = \left[\begin{array}{cc} 1 & 1 \\ 1 & 0 \end{array}\right] \tag{3.3}$$

induces the golden-mean shift Σ_A with positive entropy $h(\Sigma_A) = \log\left(\frac{1+\sqrt{5}}{2}\right)$.

To incorporate the effect of Σ_A, Table 1.1 is replaced by the following table.



Table 3.1.

As in Table 1.1, the horizontal lines in Table 3.1 connect the integers in $i\mathbb{M}_2$ for each $i \in \mathcal{I}_2$, the effect comes from $\mathbb{X}_2^0$. On the other hand, the bold zigzag line in

Table 3.1 connects all natural integers comes from Σ_A. Therefore, for any $i \neq j$ in $\mathcal{I}_2$, $i\mathbb{M}_2$ and $j\mathbb{M}_2$ are no longer mutually independent. In fact, they are all coupled through the relation set $\mathbb{M}_2$. Therefore, (3.2) is regarded as a coupled system.

Before the system $\mathbb{X}_2^A$ is decoupled, the following definition is needed.

Definition 3.1. *Two sets of integers of M and M' are mutually independent in $\mathbb{X}_2^A$ if*

$$M \cap M' = \emptyset \tag{3.4}$$

and any numbers m in M and m' in M' are not consecutive and m and m' are not consecutive in 2-power, i.e., if $m = 2^n$ for some n then $m' \neq 2^{n+1}$ and 2^{n-1}.

In this case, M and M' are decoupled in $\mathbb{X}_2^A$. The following lemma is easily obtained.

Lemma 3.2. *Suppose M and M' are mutually independent in $\mathbb{X}_2^A$. Then*

$$|\Sigma(M \cup M')| = |\Sigma(M)||\Sigma(M')|, \tag{3.5}$$

where $\Sigma(M)$ is the set of all admissible patterns on lattice M, and $\Sigma(M')$ and $\Sigma(M \cup M')$ are defined analogously.

The strategy for studying (3.2) is to decouple the whole system into disjoint pieces by eliminating $\mathbb{M}_2$, such that only

$$\tilde{\mathbb{X}}_2^A = \left(\bigcup_{1 < i \in \mathcal{I}_2} i\mathbb{M}_2 \right) \bigcap \Sigma_A \tag{3.6}$$

is considered. From the reduced system $\tilde{\mathbb{X}}_2^A$, a sequence $\left\{\mathbb{X}_2^A(m)\right\}_{m=1}^{\infty}$ of independent branches are chosen. Then, the entropy of the decoupled independent system $\mathbb{X}_2^A(m)$ can be computed easily. An appropriate choice of $\mathbb{X}_2^A(m)$ is demonstrated to enable the recovery of the entropy of $\mathbb{X}_2^A$, i.e.,

$$\lim_{m \to \infty} h(\mathbb{X}_2^A(m)) = h(\mathbb{X}_2^A). \tag{3.7}$$

As in decoupled system $\mathbb{X}_2^0$, the admissible numbered lattice M_k in $\tilde{\mathbb{X}}_2^A$ is firstly picked up. Indeed, M_k is determined as follows.



(a) M_1 (b) $M_2(3)$ (c) $M_2(l)$



(d) $M_3(3)$ (e) $M_3(l)$



(f) $M_4(3)$

Figure 3.1.

Notably, $M_m(l)$ and $M_m(l')$ are mutually independent when $l, l' \in \mathcal{I}_2$ and $|l - l'| \geq 2^{m+1}$.

Consider in more detail the construction of M_k, which is recursive and robotic. The basic concept is that any number can produce the next generation through $\mathbb{X}_2$ or Σ_A. More precisely, for each number n, if $n \in \mathcal{I}_2$, then n can produce the next generation $2n \in n\mathbb{M}_2$. If $n \notin \mathcal{I}_2$ with $n = i2^m$, $m \geq 1$, then n can produce $2n = i2^{m+1} \in i\mathbb{M}_2$ through $\mathbb{X}_2$ and $n \pm 1 = i2^m \pm 1 \in \mathcal{I}_2$ through Σ_A. A complete production cycle is as follows. If $n \in \mathcal{I}_2$, then n produces $2n$ and then $2n \pm 1$. If $n = i2^m$, $m \geq 1$, then n produces $i2^{m+1}$ and then $i2^{m+1} \pm 1$.

For example, for $M_1(3)$, only one cell exists, and number three is regarded as the number of first generation. M_2 is constructed from M_1 by extending the number three in first generation to the number six in the second generation through $3\mathbb{M}_2$, and then taking six in the first generation to create numbers five and seven as the descendanta of six through Σ_A. In the horizontal direction, M_2 has two cells that are numbered "3" and "6". M_2 is called of degree 2.

The construction of M_3 form M_2 is performed similarly: the number six yields the third-generation number 12 in $\mathbb{M}_2$. Simultaneously, numbers five and seven yield the second-generation numbers ten and 14 in $\mathbb{M}_2$, respectively. Next, numbers ten, 14 and 12 yield their descendants nine, 11, 13, 15 and 11, 13 through Σ_A, as presented in Fig 3.1 (d). The maximal number of numbers of cells in the horizontal direction is now three, and M_3 is therefore now of degree 3. On M_k, the maximal number of numbers of cells in the horizontal direction is k, and M_k is of degree k.



Figure 3.2.

After the basic lattices M_k and L_k are identified, in a given range $\mathcal{N}(2^n)$, step (II), which is the computation of the number of disjoint copies of $M_k(l) \subset \tilde{\mathbb{X}}_2^A$ with $l \in \mathcal{I}_2$, is implemented. Back to Table 3.1, $[1, 32]$ can be decoupled by $M_2(3)$, $M_2(9)$, $M_2(11)$, $M_2(13)$, $M_2(15)$ and $\{1, 2, 4, 8, 10, 12, 14, 16, 20, 24, 28, 32\}$ are not used; see Fig 3.3.



Figure 3.3.

There exist one copy $M_2(3)$ in $(2, 2^3)$ and four copies $M_2(9)$, $M_2(11)$, $M_2(13)$, $M_2(15)$ in $(2^3, 2^5)$. Notably, $[1, 2^5] \setminus \mathbb{M}_2 = M_4(3)$. However, Fig 3.3 enables us to approximate entropy by using M_2 alone under the expense of losing numbers which are relatively small compared with the numbers that have been used.

Furthermore, the following observations are made and are very useful in studying M_2 and M_3. In Fig 3.4, between $\left(2^{n-2}, 2^n\right)$, each odd integer $l \in \left(2^{n-2}, 2^{n-1}\right) \cap \mathcal{I}_2$ can produce a $M_2(l)$ that lies in $\left(2^{n-2}, 2^n\right)$. Furthermore, they are all disjoint. Therefore, there are totally 2^{n-3} copies of M_2. Similarly, as in Fig 3.5, between $\left(2^{n-3}, 2^n\right)$, each $l \in \left(2^{n-3}, 2^{n-2}\right) \cap \mathcal{I}_2$ produces a $M_3(l)$ that lies in $\left(2^{n-3}, 2^n\right)$. They are all disjoint. The total number of copies of $M_3(l)$ in $\left(2^{n-3}, 2^n\right)$ is 2^{n-4}. Therefore, the following conclusion is drawn.



Figure 3.4.



Figure 3.5.

Lemma 3.3. *For any $k \geq 1$, if $n \geq (m+1)k+1$, within $(1, 2^n)$, there are*

$$\alpha_n(k) \equiv 2^{n-k-1}\left(1+\frac{1}{2^k}+\frac{1}{2^{2k}}+\cdots+\frac{1}{2^{mk}}\right) \tag{3.8}$$

disjoint copies of $M_k(l)$ with $l \in \mathcal{I}_2$.

Finally, for step (III), denote by Σ_k the admissible patterns on L_k. Then, Lemma 3.3 yields the lower bound of the entropy $h(\mathbb{X}_2^A)$.

Lemma 3.4. *For any $k \geq 1$,*

$$\frac{1}{2\left(2^k-1\right)} log|\Sigma_k| \leq h(\mathbb{X}_2^A). \tag{3.9}$$

Now, a good upper estimate of the entropy $h(\mathbb{X}_2^A)$ remains to be found. As in the derivation of Lemma 3.3, only the number of the vertices that are not used in (3.8) and their contribution to entropy have to be determined. The following lemma is easily obtained.

Lemma 3.5. *For each $k \geq 1$, the number of the vertices of L_k satisfies*

$$\begin{aligned} |L_k| = & \quad k + 2|L_{k-1}| \\ = & \quad 2\left(2^k-1\right) - k, \end{aligned} \tag{3.10}$$

and the number of

$$\mathcal{N}(2^n) \setminus \bigcup_{l \in \mathcal{I}_2} M_k(l) \tag{3.11}$$

is

$$\frac{k}{2\left(2^k-1\right)} 2^n. \tag{3.12}$$

Moreover,

$$h(\mathbb{X}_2^A) \leq \frac{1}{2\left(2^k-1\right)} \log|\Sigma_k| + \frac{k}{2\left(2^k-1\right)} \log 2, \tag{3.13}$$

where $|\Sigma_k|$ is the number of all admissible patterns on L_k.

Proof. (3.10) is easily proved by induction. Then, (3.11) and (3.12) follow immediately. The details are omitted.

Since two possible symbols "0" and "1" can be presented on vertices in (3.11), the upper estimate (3.13) of entropy follows.

□

Lemmas 3.4 and 3.5 immediately yield the following result.

Theorem 3.6. *The entropy $h(\mathbb{X}_2^A)$ is given by*

$$h(\mathbb{X}_2^A) = \lim_{k \to \infty} \frac{1}{2\left(2^k-1\right)} \log|\Sigma_k|. \tag{3.14}$$

Furthermore,

$$\frac{1}{2\,(2^k-1)}\log|\Sigma_k| \leq h(\mathbb{X}_2^A) \leq \frac{1}{2\,(2^k-1)}\log|\Sigma_k| + \frac{k}{2\,(2^k-1)}\log 2. \tag{3.15}$$

Proof. (3.14) follows from (3.15) immediately. □

Table 3.2 presents a numerical approximation of (3.15). For $n \geq 1$, let

$$h^{(n)}(\mathbb{X}_2^A) = \tfrac{1}{2(2^n-1)}\log|\Sigma_n| \quad \text{and} \quad \bar{h}^{(n)}(\mathbb{X}_2^A) = h^{(n)}(\mathbb{X}_2^A) + \tfrac{n}{2(2^n-1)}\log 2.$$

n	2	3	4
$\lvert\Sigma_n\rvert$	9	237	213624
$h^{(n)}(\mathbb{X}_2^A)$	0.366204	0.390576	0.409066
$\bar{h}^{(n)}(\mathbb{X}_2^A)$	0.597253	0.539107	0.501485

Table 3.2.

Remark 3.7. *Whether $h(\mathbb{X}_2^A)$ can be expressed in explicit form, as is (1.4) for $h(\mathbb{X}_2^0)$ is unclear.*

Now, the result for $\mathbb{X}_2^A$ is extended to

(3.16)
$$\mathbb{X}_Q^A = \left\{(x_1,x_2,x_3,\cdots)\in\{0,1\}^{\mathbb{N}} \mid x_k x_{Qk} = 0 \text{ for all } k\geq 1 \text{ and } (x_1,x_2,\cdots)\in\Sigma_A\right\}$$

for any integer $Q \geq 3$.

Recall the main parts of the investigation of the coupled system $\mathbb{X}_2^A$ and apply them to $\mathbb{X}_Q^A$.

- $(\mathrm{I})_c$ Remove $\mathbb{M}_Q$ from $\mathbb{N} = \bigcup\limits_{i\in\mathcal{I}_Q} i\mathbb{M}_Q$ to decouple the system, yielding the reduced system $\tilde{\mathbb{X}}_Q^A = \bigcup\limits_{1<i\in\mathcal{I}_Q} i\mathbb{M}_Q \bigcap \Sigma_A$.
- $(\mathrm{II})_c$ Identify the lattice M_k (and L_k), which is the maximal connected graph of degree k. All $M_k(l)$, $l\in\mathcal{I}_Q$, are disjoint.
- $(\mathrm{III})_c$ Compute the remaining vertices in

$$\mathcal{N}(Q^n)\setminus\bigcup_{l\in\mathcal{I}_Q} M_k(l). \tag{3.17}$$

- $(\mathrm{IV})_c$ Compute the number Σ_k of admissible patterns on L_k.

Step $(\mathrm{II})_c$ will give lower bound of $h(\mathbb{X}_Q^A)$, and $(\mathrm{III})_c$ will give upper bound of $h(\mathbb{X}_Q^A)$. Then, $h(\mathbb{X}_Q^A)$ follows if the error term in $(\mathrm{III})_c$ approaches zero as n tends to infinity. To minimize the error in $(\mathrm{III})_c$, the amount of unused lattices should be as small as possible. Therefore, the choice of M_k or the graph L_k in $(\mathrm{II})_c$ should be as large as possible as far as they are decoupled.

$\mathbb{X}_3^A$ can be used to illustrate the procedures of $(\mathrm{I})_c$,$(\mathrm{II})_c$ and $(\mathrm{III})_c$. From

$$\mathbb{N} = \bigcup_{i\in\mathcal{I}_3} i\mathbb{M}_3, \tag{3.18}$$

the following figure is drawn, corresponding to Table 3.1 .



Table 3.3.

Take M_1 as being on one vertex. In Fig 3.6, a maximal connected graph M_2 with two horizontal vertices in $\bigcup_{1<i\in\mathcal{I}_3} i\mathbb{M}_3$ can be identified as follows. Notably, $M_2(4)$ and $M_2(7)$ are mutually independent.



$M_2(4)$ $M_2(7)$

Figure 3.6.

Furthermore, M_3 can be constructed from M_2 as follows.



$M_3(4)$

Figure 3.7.

The following result yields the vertices in M_k of $\tilde{\mathbb{X}}^A_Q$.

Lemma 3.8. *For any* $Q \geq 3$ *and* $k \geq 2$, *denote* $L_{Q;k}$ *be the degree* k *blank lattice. The number* $|L_{Q;k}|$ *of the vertices of* $L_{Q;k}$ *is*

$$|L_{Q;k}| = \frac{Q\left(Q^k - 1\right)}{Q-1} - k. \tag{3.19}$$

Proof. The result is proven first for the case $Q = 3$. The other cases can be treated analogously.

For $Q = 3$, let

$$a_{3,n} = 2 \cdot 3^n.$$

The blank lattice $L_{3;2}$ can be obtained from $M_2(4)$ in Fig 3.5, and

$$|L_{3;2}| = a_{3,1} + (a_{3,1} + a_{3,2}) = 10.$$

Now, $L_{3;3}$ is obtained from $M_3(4)$ in Fig 3.6 and can be grouped as follows.



Figure 3.8.

According to Fig 3.8, it is easy to see that

$$\begin{aligned} |L_{3;3}| &= \sum_{l=1}^{3}\sum_{j=1}^{l} a_{3,j} \\ &= a_{3,1} + (a_{3,1} + a_{3,2}) + (a_{3,1} + a_{3,2} + a_{3,3}) \\ &= |L_{2;2}| + 26. \end{aligned}$$

By induction, it can be proven

$$|L_{3;m}| = |L_{3;m-1}| + 3^m - 1. \tag{3.20}$$

Therefore, (3.19) follows for $Q = 3$.

For $Q = 4$, $M_{4,2}$ is as in Fig 3.9.



$M_{4;2}$

Figure 3.9.

Similarly, for any $Q \geq 4$, $M_{Q,2}$ is given in Fig 3.10.



$c_1 = Q^2+1$, $c_2 = Q^2+Q-1$, $c_3 = Q^2+Q+1$, $c_4 = Q^2+2Q-1$, $c_5 = Q^2+Q+1$,
$c_6 = Q^2+3Q-1$, $c_7 = Q^2+3Q+1$, $c_8 = 2Q^2-Q-1$, $c_9 = 2Q^2-Q+1$, $c_{10} = 2Q^2+2Q-1$

$M_{Q,2}$

Figure 3.10.

Define

$$a_{Q,n} = (Q-1)2^{n-1}.$$

Careful mathematical induction proves (3.19). The details are omitted here. □

Now, we can extend Theorem 3.6 to all $Q \geq 3$, as follows.

Theorem 3.9. *For any $Q \geq 3$ and $k \geq 2$,*

$$\frac{Q-1}{Q\left(Q^k-1\right)} \log|\Sigma_{Q;k}| \leq h(\mathbb{X}_Q^A) \leq \frac{Q-1}{Q\left(Q^k-1\right)} \left(\log|\Sigma_{Q;k}| + k\log 2\right), \tag{3.21}$$

and

$$h(\mathbb{X}_Q^A) = \lim_{k\to\infty} \frac{Q-1}{Q\left(Q^k-1\right)} \log|\Sigma_{Q;k}|, \tag{3.22}$$

where $\Sigma_{Q;k}$ is the set of all admissible patterns on $L_{Q;k}$.

Proof. It can be verified that the number of $M_{Q;k}(l)$, $l \in \mathcal{I}_Q \cap [1, Q^n]$, is

$$\frac{Q-1}{Q\left(Q^k-1\right)} Q^{n-1}. \tag{3.23}$$

By (3.19), the number of the remaining vertices in $[1, Q^n]$ is

$$\frac{(Q-1)k}{Q\left(Q^k-1\right)}.$$

The results follow immediately.

□

Like Theroem 2.12, Theorem 3.9 can be generalized to any number of symbols, any constraints $\mathcal{C}$ and any additive shifts of finite type Σ_A. Indeed, let

$$\begin{aligned}&\mathbb{X}_Q^A(N,\mathcal{C}) \\ =\ &\left\{(x_1,x_2,x_3,\cdots)\in\{0,1,\cdots,N-1\}^{\mathbb{N}} \mid x_k x_{Qk}\in\mathcal{C} \text{ for all } k\geq 1 \text{ and } (x_1,x_2,x_3,\cdots)\in\Sigma_A\right\},\end{aligned} \tag{3.24}$$

where $\mathcal{C}\subseteq\left\{0,1,\cdots,(N-1)^d\right\}$ is a constraint set and A is an $m\times m$ $0-1$ matrix.

Then, the following theorem can easily be obtained. The details of the proof are omitted.

Theorem 3.10. *For any $Q\geq 3$, $\mathcal{C}\subseteq\left\{0,1,\cdots,(N-1)^d\right\}$ and $k\geq 2$,*

$$\frac{Q-1}{Q\left(Q^k-1\right)}\log|\Sigma_k(Q;A;N,\mathcal{C})|\leq h\left(\mathbb{X}_Q^A(N,\mathcal{C})\right)\leq\frac{Q-1}{Q\left(Q^k-1\right)}\left(\log|\Sigma_k(Q;A;N,\mathcal{C})|+k\log N\right), \tag{3.25}$$

and

$$h\left(\mathbb{X}_Q^A(N,\mathcal{C})\right)=\lim_{k\to\infty}\frac{Q-1}{Q\left(Q^k-1\right)}\log|\Sigma_k(Q;A;N,\mathcal{C})|, \tag{3.26}$$

where $\Sigma_k(Q;A;N,\mathcal{C})$ is the set of all admissible patterns on $L_{Q;k}$, the constraint of the vertices on the bold lines in $L_{Q;k}$ is given by A and the constraint of the vertices on thin lines in $L_{Q;k}$ is given by N and $\mathcal{C}$.

Acknowledgement. The numerical results of this paper are provided by Mr. Hung-Shiun Chen, a Ph.D. student of Song-Sun Lin at National Chiao Tung University.

References

1. J.C. Ban, C.H. Chang and S.S. Lin, *On The Structure of Multi-Layer Cellular Neural Networks*, J. Differential Equations, **252**, no.8, (2012) 4563–4597.
2. J.C. Ban, C.H. Chang, S.S. Lin and Y.H. Lin, *Spatial complexity in multi-layer cellular neural networks*, J. Differential Equations, **246**, (2009) 552–580.
3. J.C. Ban, W.G. Hu, S.S. Lin and Y.H. Lin, *Zeta functions for two-dimensional shifts of finite type*, to appear in Memo. Amer. Math. Soc..
4. J.C. Ban, W.G. Hu, S.S. Lin and Y.H. Lin, *Verification of mixing properties in two-dimensional shifts of finite type*, Preprint arXiv:1112.2471.
5. J.C. Ban and S.S. Lin, *Patterns generation and transition matrices in multi-dimensional lattice models*, Discrete Contin. Dyn. Syst. **13**, no. 3, (2005) 637–658.
6. J.C. Ban, S.S. Lin and Y.H. Lin, *Patterns generation and spatial entropy in two dimensional lattice models*, Asian J. Math. **11**, no. 3, (2007) 497–534.
7. J.C. Ban, S.S. Lin and Y.H. Lin, *Three-dimensional cellular neural networks and pattern generation problems*, International J. Bifurcation and Chaos. **18**, no. 4, (2008) 957–984.
8. A. Fan, L. Liao, J. Ma, *Level sets of multiple ergodic averages*, Preprint arXiv:1105.3032, to appear in Monatsh. Math.
9. A. Fan, J. Schmeling, M.Wu, *Multifractal analysis of multiple ergodic averages*, C. R. Math. Acad. Sci. Paris 349 (2011), 961-964.
10. B. Grünbaum and G. C. Shephard, *Tilings and Patterns*, New York: W. H. Freeman, (1986).
11. W.G. Hu and S.S. Lin, *Zeta functions for higher-dimensional shifts of finite type*, International J. of Bifurcation and Chaos, **19**, no. 11, (2009) 3671–3689.

12. W.G. Hu and S.S. Lin, *Nonemptiness problems of plane square tiling with two colors*, Proc. Amer. Math. Soc., **139**, (2011) 1045–1059.
13. J. Juang and S.S. Lin, *Cellular Neural Networks: Mosaic pattern and spatial chaos*, SIAM J. Appl. Math, **60**, no. 3, (2000) 891–915.
14. R. Kenyon, Y. Peres and B. Solomyak, *Hausdorff dimension for fractals invariant under the multiplicative integers*, Ergodic Theory Dynam. Sys., to appear.
15. S.S. Lin and T.S. Yang, *On the spatial entropy and patterns of two-dimensional cellular neural networks*, International J. of Bifurcation and Chaos. **12**, (2002) 115–128.
16. D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press, Cambridge, (1995).
17. Y. Peres, J. Schmeling, S. Seuret and B. Solomyak, *Dimensions of some fractals defined via the semigrop generated by* 2 *and* 3, Preprint arXiv:1206.4742.
18. Y. Peres, B. Solomyak, *Dimension spectrum for a nonconventional ergodic average*, Preprint arXiv:1107.1749, to appear in Real Analysis Exchange.
19. N. J. A. Sloane and S. Plouffe, *The encyclopedia of integer sequences*, Academic Press, 1995.

Department of Applied Mathematics, National Dong Hwa University, Hualien 97401, Taiwan
E-mail address: `jcban@mail.ndhu.edu.tw`

Department of Applied Mathematics, National Chiao Tung University, Hsin-Chu 30010, Taiwan
E-mail address: `wghu@mail.nctu.edu.tw`

Department of Applied Mathematics, National Chiao Tung University, Hsin-Chu 30010, Taiwan
E-mail address: `sslin@math.nctu.edu.tw`